

Sugama, H.

Horton, W. (Inst. Fusion Studies, Univ. Texas)

Neoclassical transport processes of electrons and ions are investigated in detail for toroidally rotating axisymmetric plasmas with large flow velocities on the order of the ion thermal speed [1]. The magnetic field and the toroidal flow velocity $\mathbf{V}_0 = \mathcal{O}(v_{Ti})$ are written as $\mathbf{B} = I\nabla\zeta + \nabla\zeta \times \nabla\Psi$ and $\mathbf{V}_0 = -Rc(\partial\Phi_0/\partial\Psi)\hat{\zeta}$, where ζ is the toroidal angle, Ψ represents the poloidal flux, and Φ_0 denotes the lowest-order electrostatic potential in the small-gyroradius expansion. For the rotating plasma consisting of electrons and single-species ions, the neoclassical transport equations are written as

$$\begin{bmatrix} \Gamma_e \\ q_e/T_e \\ q_i/T_i \\ \Pi_i \\ J_E \end{bmatrix} = \begin{bmatrix} L_{11}^{ee} & L_{12}^{ee} & L_{12}^{ei} & L_{1V}^e & L_{1E}^e \\ L_{21}^{ee} & L_{22}^{ee} & L_{22}^{ei} & L_{2V}^e & L_{2E}^e \\ L_{21}^{ie} & L_{22}^{ie} & L_{22}^{ii} & L_{2V}^i & L_{2E}^i \\ L_{V1}^e & L_{V2}^e & L_{V2}^i & L_{VV} & L_{VE} \\ L_{E1}^e & L_{E2}^e & L_{E2}^i & L_{EV} & L_{EE} \end{bmatrix} \begin{bmatrix} X_{e1}^* \\ X_{e2} \\ X_{i2} \\ X_V \\ X_E \end{bmatrix}.$$

Here, the neoclassical radial particle flux, electron and ion heat fluxes are denoted by $\Gamma_e (= \Gamma_i)$, q_e , and q_i , respectively. Π_i represents the neoclassical radial flux of the toroidal momentum, and $J_E \equiv \langle BJ_{\parallel} \rangle / \langle B^2 \rangle^{1/2}$ is the averaged parallel current. The thermodynamic forces are given by $X_{e1}^* \equiv -N_e^{-1} \partial(N_e T_e) / \partial r - N_i^{-1} \partial(N_i T_i) / \partial r$, $X_{e2} \equiv -\partial T_e / \partial r$, $X_{i2} \equiv -\partial T_i / \partial r$, $X_V \equiv -\partial(V_0/R) / \partial r$, and $X_E \equiv \langle BE_{\parallel} \rangle / \langle B^2 \rangle^{1/2}$.

The neoclassical transport coefficients L 's depend on the toroidal flow velocity V_0 . The Onsager relations for the flow-dependent neoclassical transport coefficients are derived from the symmetry properties of the drift kinetic equation with the self-adjoint collision operator. The complete neoclassical transport matrix with the Onsager symmetry is obtained for the Pfirsch-Schlüter and banana regimes.

The toroidal flow dependence of the ion thermal diffusivity $\chi_i = T_i L_{22}^{ii} / n_i$ is expressed in

terms of the enhancement factor $F(\Upsilon)$ which is given for the banana regime as

$$\begin{aligned} F(\Upsilon) &\equiv \frac{\chi_i(V_0)}{\chi_i(V_0=0)} \\ &= 1 + 0.765\Upsilon - 0.631\Upsilon^2 + 0.280\Upsilon^3 \end{aligned}$$

where

$$\Upsilon \equiv \frac{m_i V_0^2}{(T_e + T_i)}.$$

This enhancement factor is in good agreement with that given by Catto *et al.* in spite of the difference between the solution methods.

The full expression of the parallel current for the toroidally rotating plasma in the banana regime is given by

$$\begin{aligned} J_E = & - \left(\frac{r}{R_0} \right)^{1/2} \frac{c}{B_P} \left[2.411 F_{E1}^e(\Upsilon) \frac{dP}{dr} \right. \\ & \left. - 1.800 F_{E2}^e(\Upsilon) n_e \frac{dT_e}{dr} - 2.828 F_{E2}^i(\Upsilon) n_i \frac{dT_i}{dr} \right] \\ & + \sigma_S \left[1 - 1.832 \left(\frac{r}{R_0} \right)^{1/2} F_{EE}(\Upsilon) \right] X_E \end{aligned}$$

with the Spitzer conductivity σ_S and the enhancement factors

$$\begin{aligned} F_{E1}^e(\Upsilon) &= 1 + 0.868\Upsilon - 0.539\Upsilon^2 + 0.229\Upsilon^3 \\ F_{E2}^e(\Upsilon) &= 1 + 2.248\Upsilon - 1.661\Upsilon^2 + 0.727\Upsilon^3 \\ F_{E2}^i(\Upsilon) &= 1 + 1.494\Upsilon - 1.022\Upsilon^2 + 0.434\Upsilon^3 \\ F_{EE}(\Upsilon) &= 1 + 0.431\Upsilon - 0.184\Upsilon^2 + 0.072\Upsilon^3. \end{aligned}$$

The sum of the neoclassical and classical toroidal momentum diffusivity for the banana regime is written as $\chi_\phi = L_{VV} / (n_i m_i R_0^2) + \frac{3}{5} \rho_i^2 / \tau_i = (\frac{1}{10} q^2 + \frac{3}{5}) \rho_i^2 / \tau_i$. The banana inward-pinch velocity for the particles and the toroidal momentum due to the parallel electric field is derived from $v_{\text{inward}} = -L_{1E}^e X_E / n_e = -L_{VE}^e X_E / (n_i m_i R_0 V_0)$ as

$$v_{\text{inward}} = 2.411 F_{E1}^e(\Upsilon) \left(\frac{r}{R_0} \right)^{1/2} \frac{c E_{\parallel}}{B_P}.$$

References

- 1) Sugama, H. and Horton, W. : Phys. Plasmas **4** (1997) 2215.